

AD-A129 076

ALL SPECTRAL DOMINANT NORMS ARE STABLE(U) WISCONSIN
UNIV-MADISON MATHEMATICS RESEARCH CENTER

1/1

S FRIEDLAND ET AL. APR 83 MRC-TSR-2507 DAAG29-80-C-0041

UNCLASSIFIED

F/G 12/1

NL



END
DATA
FORMED
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A129076

THE UNIVERSITY OF MICHIGAN LIBRARY

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

DATE 11-11-83 BY 1045

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

DTIC FILE COPY

April 1983

(Received April 6, 1983)

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

DTIC
ELECTE
JUN 9 1983

S

A

88 06 07 074

UNIVERSITY OF SICHUAN-CHANGSHU
MATHEMATICS RESEARCH CENTER

ALL SPECTRAL DOMINANT NORMS ARE STABLE

S. Friedland^{*} and C. Sogor^{**}

Technical Summary Report 02507
April 1983

ABSTRACT

A vector norm $|\cdot|$ on the space of $n \times n$ complex valued matrices is called stable if

$$|A^m| \leq K|A|^m$$

for all A and non-negative integers m . We show that such a norm is stable if and only if it dominates the spectral radius.

AMS (MOS) Subject Classifications: 15A60, 39A11

Key Words: Stable norms, Numerical ranges

Work Unit Number 3 (Numerical Analysis and Scientific Computing)

^{*}Hebrew University at Jerusalem.

^{**}Technische Universitat, Munchen.

SIGNIFICANCE AND EXPLANATION

When solving partial differential equations numerically one often has to use iterations involving matrices restricted to a given set A of $n \times n$ complex valued matrices. It then follows that the iteration scheme is stable if and only if this set of matrices is stable. That is all powers of all matrices from the set A are uniformly bounded. Such sets were completely characterized by H. O. Kreiss. However, his criteria are hard to use.

the authors
In this paper we characterize in a very simple way stable sets of matrices A , whenever the set A is closed, convex, balanced, and contains a neighborhood of the origin. Such a set A is a unit ball of some vector norm $\|\cdot\|$ on matrices. *They* We then show that A is stable if and only if the above norm dominates the spectral radius ρ . That is $\rho(A) \leq \|A\|$ for all matrices A . The necessity of the above condition is obvious, and is sometime referred to as the Neumann condition. *the authors* To prove the sufficiency we use the Kreiss matrix theorem and other results.

TRUE VALUE OF .



By	
Distribution	
Availability Codes	
Dist	Special
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

ALL SPECTRAL CONSTANT SETS ARE STABLE

S. Friedland* and C. Berger**

1. Introduction

Let A be a set of $n \times n$ complex valued matrices - $M_n(C)$. A is called stable if

$$(1.1) \quad \|A^m\| < K, \quad m = 0, 1, 2, \dots, \quad A \in A.$$

Here $\|\cdot\|$ is a vector norm on $M_n(C)$.

In 1962 Kreiss [5] characterized stable sets. In particular he showed that (1.1) is equivalent to

$$(1.2) \quad \|(zI - A)^{-1}\| < C/(|z| - 1), \quad \text{for all } |z| > 1, \quad A \in A.$$

While (1.1) easily implies (1.2) with $K = C$ it can be shown that (1.2) implies (1.1) with

$$(1.3) \quad K = \alpha_n C, \quad \alpha_n \leq \frac{32en}{n}, \quad \lim_{n \rightarrow \infty} \alpha_n = 0.$$

See for example [7] and [9].

The serious draw-back of (1.2) is that it is difficult to verify in general. Thus, a natural question is whether the condition (1.2) can be replaced by a simpler condition assuming the set A is of a certain type. In many instances A is of the following type:

- (i) A - is closed,
- (ii) A - is convex,
- (iii) A - is circular, i.e. $e^{i\theta} A = A$ for all $\theta \in R$,
- (iv) A - contains an open set.

* Hebrew University at Jerusalem.

** Technische Universität, München.

Clearly, these conditions are equivalent to the assumption that A is a unit ball of some vector norm $|\cdot|$ on $M_n(C)$.

$$(1.4) \quad A = \{A, |A| \leq 1\}.$$

Thus, $|\cdot|$ is called stable if its unit ball is a stable set. For $A \in M_n$, let $\rho(A)$ denote the spectral radius of A . Since on finite dimensional vector space all the norms are equivalent we have the equality

$$(1.5) \quad \rho(A) = \lim_{m \rightarrow \infty} |A^m|^{1/m}.$$

So, if A is a stable set we get

$$(1.6) \quad \rho(A) < 1, \quad A \in A.$$

Thus if $|\cdot|$ is a stable norm we have that $\rho(A) < 1$ for $|A| = 1$. Using the homogeneity of $\rho(\cdot)$ and $|\cdot|$ we get

$$(1.7) \quad \rho(A) < |A|.$$

Recall that $|\cdot|$ is called spectrally dominant if (1.7) holds. Our main result is

Theorem 1. Let $|\cdot|$ be a vector norm on $M_n(C)$. Then $|\cdot|$ is stable if and only if it is spectrally dominant.

This result was conjectured by C. Johnson in [4]. The case of unitary invariant norms was proved in Friedland-Tadmor in [3].

3. Main Results

Following Benger [10] we first consider special spectral dominant norms on $M_n(C)$. These norms are called the generalized numerical radius and are denoted by $r_g(\cdot)$. For reader's convenience we give short proofs of these known results. Let $\|\cdot\|_2$ be the standard Euclidean norm on C^n . As usual let x be a column vector in C^n , x^t and x^* its transpose and conjugate transpose. Denote by S_2 the unit sphere of this norm.

Assume that we have the following map

$$(2.1) \quad \phi : S_2 \rightarrow 2C^n.$$

We suppose that

$$(2.2) \quad y^t x = 1, \text{ for all } y \in \phi(x).$$

We now assume that the map (2.1) is closed. That is, if $x_k \in S_2$, $y_k \in \phi(x_k)$, $x_k \rightarrow x$, the sequence $\{y_k\}$ is bounded. Moreover if $y_k \rightarrow y$ then $y \in \phi(x)$. This in particular implies that $\phi(x)$ is compact and $\bigcup_{x \in S_2} \phi(x)$ is bounded. We then define the generalized numerical radius as

$$(2.3) \quad r_g(A) = \max_{x \in S_2, y \in \phi(x)} |y^t A x|.$$

Lemma 1. The generalized numerical range is a spectral dominant norm on $M_n(C)$.

Proof: In view of (2.2) - (2.3) we have that

$$(2.4) \quad \rho(A) \leq r_g(A).$$

Also (2.3) yields that $r_g(A)$ is a seminorm. Assume that $r_g(A) = 0$ and $A \neq 0$.

According to (2.4) A is nilpotent. Choose a basis in C^n such that A is of the form

$$A = \left(\begin{array}{cccc|c} 0 & 1 & \dots & 0 & \\ & \cdot & & \cdot & \\ & & \cdot & \cdot & \\ & & & 1 & \\ 0 & & & & 0 \\ \hline & & & & A_1 \end{array} \right).$$

Let B be of the form

$$B = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ \hline & \\ & 0 \end{pmatrix}.$$

Then for $\epsilon > 0$

$$\rho(A + \epsilon B) = \sqrt{\epsilon} < r_g(A + \epsilon B) = \epsilon r_g(B).$$

Clearly this inequality can not hold for any $\epsilon > 0$. The above contradiction shows that $r_g(A)$ is a vector norm.

Let $\|\cdot\|$ be a norm on $X = \mathbb{C}^n$. Denote by $\|\cdot\|^*$ the corresponding norm of the dual space X^* ,

$$\|y\|^* = \max_{\|x\| \leq 1} |y^t x|.$$

We then let

$$\phi(x) = \{y, y^t x = 1, \|y\|^* \|x\| = 1\}, \quad (x \neq 0).$$

It is easy to show that in this case the map (2.1) is closed. The corresponding generalized numerical radius is the Bauer numerical radius with respect to the norm $\|\cdot\|$. See [2] for details and references.

Denote by U the set of unitary matrices in $M_n(\mathbb{C})$. As usual by $\|\cdot\|_2$ we denote the induced operator norm on $M_n(\mathbb{C})$

$$(2.5) \quad \|A\|_2 = \max_{\|x\|_2 \leq 1} \|Ax\|_2.$$

Theorem 2. Let $r_g(\cdot)$ be a generalized numerical radius. Put

$$(2.6) \quad C = \max_{U \in U} r_g(U).$$

Then

$$(2.7) \quad \|(zI - A)^{-1}\|_2 < C/(|z| - 1) \text{ for all } |z| > 1, r_g(A) < 1.$$

In particular a generalized numerical radius is a stable norm.

Proof: We first note that

$$(2.8) \quad \|y\|_2 < C \text{ for all } y \in \phi(x).$$

Indeed for $u \in E_2$ there exists $0 < \alpha < 1$ such that $r^2 u = \alpha u$. Then (2.6) follows from (2.5). Assume next that $r_q(A) < 1$. So $(zI - A)^{-1}$ is defined for $|z| > 1$. Let

$$v = (zI - A)^{-1}u, u \in E_2, v_1 = v/v_2.$$

Then, for $y \in \theta(v_1)$ we have

$$\frac{|y^t u|}{|v|_2} = |y^t (zI - A)v_1| = |z - y^t A v_1| > |z| - 1.$$

On the other hand

$$|y^t u| \leq |y|_2 |u|_2 \leq C.$$

Combine the above inequalities to get

$$|(zI - A)^{-1}u|_2 \leq C/(|z| - 1), \quad |u|_2 = 1.$$

This proves (2.7). Now the stability of the generalized numerical radius follows from the Kreiss matrix theorem.

Finally Theorem 1 follows from Theorem 2 and Zenger's theorem [10] whose proof we bring for reader's convenience.

Theorem 3. (Zenger). Let $|\cdot|$ be a spectral dominant norm on $M_n(C)$. Then there exist a generalized numerical radius $r_y(\cdot)$ which is subordinate to $|\cdot|$. That is

$$(2.9) \quad r_y(A) \leq |A| \text{ for all } A \in M_n(C).$$

Proof: Let A be the unit ball of $|\cdot|$. Consider the convex balanced set

$$A_1 = \{B, B = (1 - \alpha)A + zI, z \in C, 0 < \alpha < 1, |z| < \alpha\}.$$

Clearly

$$\sigma(B) = (1 - \alpha)\sigma(A) + z$$

where $\sigma(B)$ is the spectrum of B . As $\rho(A) < 1$ for $A \in A$ we have that

$$(2.10) \quad \rho(B) < 1 \text{ for } B \in A_1.$$

Also as $A \subseteq A_1$, A_1 is the unit ball of a new norm $|\cdot|_1$ such that

$$(2.11) \quad |A|_1 \leq |A|.$$

The inequality (2.10) implies that $|\cdot|_1$ is also spectral dominant. For $|u|_2 = 1$ let

$$(2.12) \quad B(x) = \{u, u = Ax, |A|_1 < 1\}.$$

Clearly $\phi(x)$ is convex and $x \in \phi(x)$. The separation theorem yields the existence of $y \in \mathbb{C}^n$ such that

$$\operatorname{Re}(y^*x) > \operatorname{Re}(y^*Ax), \quad |A|_1 < 1.$$

Since $|a|_1 < 1$ for $|a| < 1$ we immediately deduce that y^*x is real and positive.

Thus we can normalize y such that $y^*x = 1$. Let $\phi(x)$ be the set of all u such that

$$(2.13) \quad u^*x = 1, \quad \max_{|A|_1 < 1} |u^*Ax| = 1.$$

Clearly, $y \in \phi(x)$. As

$$(2.14) \quad u^*Ax = \operatorname{tr}(Axu^*),$$

we deduce that

$$(2.15) \quad |xu^*|_1^* = 1$$

where $|\cdot|_1^*$ is the conjugate norm to $|\cdot|_1$ on $M_n(\mathbb{C})$

$$(2.16) \quad |B|_1^* = \max_{|A|_1 < 1} |\operatorname{tr}(AB)|.$$

This in particular implies that the map ϕ is closed. Since any finite dimensional vector space is reflexive we have that

$$(2.17) \quad |A|_1 = \max_{|B|_1^* < 1} |\operatorname{tr}(AB)|.$$

Compare (2.3), (2.14), (2.15) with (2.16) to deduce

$$(2.18) \quad r_g(A) < |A|_1.$$

Then (2.11) implies (2.9).

1. Stable Norms

Let $r(A)$ be the standard numerical radius of A

$$(3.1) \quad r(A) = \max_{\|x\|=1} |\langle Ax, x \rangle|.$$

The Schwarz inequality yields

$$(3.2) \quad r(A^m) \leq r(A)^m.$$

See for example [8] for a short proof.

We call the unit ball A of a vector norm to be super stable, if

$$(3.3) \quad A^m \subset A, \quad m = 1, 2, \dots$$

where

$$(3.4) \quad A^m = \{B, B = A^m, A \in A\}.$$

The inequality (3.2) implies that $r(\cdot)$ is a super stable norm. This in particular yields the Lax-Wendroff result [6] that the set $r(A) < 1$ is stable. In fact, Theorem 1 is a natural extension of the Lax-Wendroff condition.

Problem 1. Characterize all spectral dominant norms on $M_n(C)$ which are super stable.

Clearly, the standard operator norm on $M_n(C)$ is super stable.

In many numerical schemes for solutions of partial equations one has to consider a stable set of matrices A whose order is not fixed, but in fact can be arbitrary large. In that case the Kreiss matrix theorem does not apply. See for example [7]. Therefore one needs to study stable sets in the infinite dimensional case. Let B be a Banach space with a norm $\|\cdot\|$, $L(B)$ the space of all linear bounded operators $T : B \rightarrow B$ with the induced operator norm $\|\cdot\|$. Assume that $|\cdot|$ is a norm on $L(B)$ which is equivalent to the operator norm

$$(3.5) \quad \alpha \|T\| \leq |T| \leq \beta \|T\|, \quad 0 < \alpha < \beta.$$

As before, we call $|\cdot|$ stable if the unit ball of this norm is stable, i.e., (1.1) holds.

and refer to the eigenvalues of A by

$$(3.6) \quad \lambda_j(A) = \lambda_j(A) \quad j=1, \dots, n$$

the inequality (3.6) yields that a stable norm is spectral dominant

Problem 1. Characterize all norms on \mathbb{C}^n which are stable and equivalent to the spectral norm.

We note that there are spectral dominant norms on \mathbb{C}^n which are not stable and are equivalent to the operator norm. Indeed, if we choose $|B|$ to be the numerical radius of B with respect to the given norm (1.1) we then have the inequalities

$$|B| < \|B\| < \alpha |B|.$$

See for example [2]. Furthermore, according to Hille's [1], Theorem 2, there exists an operator B such that $|B| = 1$, $\|B\| = \alpha$, $\|B^k\| > \sqrt{k}$, $k = 2, 3, \dots$.

Finally we close our paper with a very specific problem. For $x = (x_1, \dots, x_n)^t \in \mathbb{C}^n$ let

$$(3.7) \quad \|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}, \quad 1 < p < \infty$$

$$y(x) = (\bar{x}_1 |x_1|^{p-2}, \dots, \bar{x}_n |x_n|^{p-2})^t, \quad x \neq 0.$$

Then, for $A \in M_n(\mathbb{C})$ we define $r_p(A)$ - the p -th numerical radius

$$(3.8) \quad r_p(A) = \max_{\|x\|_p=1} |y^t(x)Ax|.$$

Theorem 1, in this case is equivalent to the inequality

$$(3.9) \quad r_p(A^m) \leq K_{p,n} r_p(A)^m, \quad m = 0, 1, 2, \dots,$$

for all $A \in M_n(\mathbb{C})$. The inequality (3.2) yields that

$$(3.10) \quad K_{2,n} = 1.$$

We may assume in (3.9) that $K_{p,n}$ is best possible. In that case clearly

$$(3.11) \quad K_{p,n} \leq K_{p,n+1}.$$

Let q be conjugate to p

$$(3.12) \quad p^{-1} + q^{-1} = 1.$$

Since $\| \cdot \|_p$ is the conjugate norm to $\| \cdot \|_q$ on \mathbb{R}^n

we have

$$\| \cdot \|_p = \left(\sum_{j=1}^n |x_j|^p \right)^{1/p}$$

Therefore

(3.10)

$$\| \cdot \|_p = \left(\sum_{j=1}^n |x_j|^p \right)^{1/p}$$

LEMMA 3. For each value of $p \in [1, \infty)$ the sequence $\{K_{p,n}\}_{n=1}^{\infty}$ is bounded

in the case the matrix norm $p = 1, \infty$ it is clear that one can define $K_{p,n}$ as

a scalar for $p = 1, \infty$. Also we can let

(3.11)

$$r_1(A) = \lim_{p \rightarrow 1} r_p(A), \quad r_{\infty}(A) = \lim_{p \rightarrow \infty} r_p(A)$$

Let $\|A\|_p$ be the induced operator norm on $\mathbb{R}^n(\mathbb{C})$. Then it is easy to show that

(3.12)

$$r_1(A) = \|A\|_1, \quad r_{\infty}(A) = \|A\|_{\infty}$$

So

(3.17)

$$K_{1,n} = K_{\infty,n} = 1$$

The equalities (3.10) and (3.17) suggest that $\{K_{p,n}\}_{n=1}^{\infty}$ is always bounded.

REFERENCES

- [1] B. Bollobás, The power inequality on Banach spaces, Proc. Camb. Soc. 69 (1971), 411-415.
- [2] F. F. Bonsall and J. Duncan, Numerical Ranges I and II, Cambridge University Press 1971 and 1973, London.
- [3] S. Friedland and E. Tadmor, Optimality of the Lax-Wendroff condition, Linear Algebra and Appl., to appear.
- [4] C. R. Johnson, Power inequalities and spectral dominance of generalized matrix norms, Linear Algebra and Appl. 28 (1979), 117-130.
- [5] H-O. Kreiss, Über die Stabilitätsdefinition für Differenzengleichungen die Differentialgleichungen approximieren, BIT 2 (1962), 153-181.
- [6] P. D. Lax and B. Wendroff, Difference schemes for hyperbolic equations with high order of accuracy, Comm. Pure Appl. Math. 17 (1964), 384-398.
- [7] C. A. McCarthy and J. Schwartz, On the norm of a finite Boolean algebra of projections and applications to theorems of Kreiss and Morton, Comm. Pure Appl. Math. 18 (1965), 191-201.
- [8] C. Pearcy, An elementary proof of the power inequality for the numerical radius, Mich. Math. J. 13 (1966), 289-291.
- [9] E. Tadmor, The equivalence of L_2 -stability, the resolvent condition, and strict H -stability, Linear Algebra and Appl. 41 (1981), 151-159.
- [10] C. Zenger, Minimal subadditive inclusion domains for the eigenvalues of matrices, Linear Algebra and Appl. 17 (1977), 233-268.

SF/CZ/scr

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2507	2. GOVT ACCESSION NO. AD-A129076	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ALL SPECTRAL DOMINANT NORMS ARE STABLE		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) S. Friedland and C. Zenger		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis and Scientific Computing
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE April 1983
		13. NUMBER OF PAGES 10
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Stable norms Numerical ranges		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A vector norm $ \cdot $ on the space of $n \times n$ complex valued matrices is called stable if $ A^m \leq K A ^m$ for all A and non-negative integers m . We show that such a norm is stable if and only if it dominates the spectral radius.		

